

# Twisted light could dramatically boost internet speeds

April 8 2016, by Min Gu And Qiming Zhang, Rmit University

---



Credit: AI-generated image ([disclaimer](#))

[Fibre optics](#) allow for the communication of data at the speed of light.

But the amount of data that can be sent along any optic fibre is limited by how much [information](#) you can encode into the light wave travelling through it.

Currently, optic fibre technology uses several different properties of light to encode information, including brightness, colour, polarisation and direction of propagation.

But if we want to cram even more information through optic fibre, we need to use other features of light to encode more information, without disrupting currently used properties.

Such a feature could help boost the bandwidth of optic fibre technology, including our internet speeds.

## Detecting the twist

If the [light wave](#) travelling through the optic fibre is twisted helically – like a spring – then it has [angular momentum](#), which is a measure of its momentum when it rotates around a point.

But there was a major problem with using angular momentum to decode the information from the optic fibre. We needed a material with tiny nanoscale helical structures that could detect the [twisted light](#).

Our research, published today in [Science](#), [shows how we can control](#) the angular momentum of light at a nanoscale using an integrated photonic chip.


So for the first time, we have a chip with a series of elaborate nano-apertures and nano-grooves that allow for the on-chip manipulation of twisted light.

The helical design of these tiny apertures and grooves removes the need for any other bulky interference-based optics to detect the angular momentum signals.

So if you send an optical data signal to a [photonic chip](#), which is a microchip that uses light instead of electrons, then it is important to know where the data is going, otherwise information will be lost.

Using our nanophotonic chip, we can precisely guide angular momentum data signals without losing the information they carry.

What's more, the angular momentum information of many different signals can be processed at the same time through the chip.



The nanophotonic chip magnified 2,000 times. Each indentation on the image is a single unit of the chip, like a single pixel in a display panel, made up of semi-circle nano-grooves and nano-apertures engraved in a metallic film. Credit: RMIT University, Author provided

This means we can potentially achieve an ultra-wide bandwidth, with six-


orders magnitude of increased data access compared to current technology.

## **Technology for today**

Owing to the rapid development of nano-fabrication technology, we believe there is no technical challenge to the mass production of this chip today.

This breakthrough opens an entirely new perspective in employing [light](#) for chip-scale information generation, transmission and retrieval of images, videos, sounds and so on.

It could be used in applications such as data transmission, ultra-high definition displays, ultra-high capacity optical communications and ultra-secure optical encryption.



Professor Min Gu with the nanophotonic chip that can harness the angular momentum of light. Credit: RMIT University, Author provided

For example, the communication speed on the National Broadband Network can be boosted through the parallel processing of the angular momentum.

Since the chip consists of an array of individually-controlled single units, and each single unit is capable of independently processing the angular momentum information, this chip device allows for parallel processing of optical information.

A large number of optical fibres in one fibre bundle can be processed through the chip in parallel, which means the processing speed can be

significantly increased by considering how large the array is.

For example, if we take 100 by 100 of such units in the array for the [chip](#), then the speed could be boosted by four orders of magnitude.

This quirk of physics could one day lead to significantly faster internet speeds along with a host of other useful applications.

**More information:** "On-chip noninterference angular momentum multiplexing of broadband light" [DOI: 10.1126/science.aaf1112](https://doi.org/10.1126/science.aaf1112)

*This article was originally published on [The Conversation](#). Read the [original article](#).*

Source: The Conversation

Citation: Twisted light could dramatically boost internet speeds (2016, April 8) retrieved 15 April 2024 from <https://techxplore.com/news/2016-04-boost-internet.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--