

Gripping geckos' aerial escapes test their limits

July 19 2017, by Krishna Ramanujan



A Madagascar day gecko in the Antwerp Zoo. Credit: Frank Wouters, Antwerpen, Belgium (Shining Star)

Geckos climb vertically up trees, walls and even windows, thanks to pads

on the digits of their feet that employ a huge number of tiny bristles and hooks.

Scientists have long marveled at the gecko's adhesive capabilities, which have been described as 100 times more than what is needed to support their body weight or run quickly up a [surface](#).

But a new theoretical study examines for the first time the limits of [geckos'](#) gripping ability in natural contexts. The study, published July 10 in the *Journal of the Royal Society Interface*, reports there are circumstances – such as when geckos fear for their lives, leap into the air and are forced to grab on to a leaf below – when they need every bit of that fabled adhesive ability, and sometimes it's not enough.

"Geckos are notoriously described as having incredible ability to adhere to a surface," said Karl Niklas, professor of plant evolution in Cornell's Plant Biology Section in the School of Integrative Plant Science and a co-author of the paper. The study's lead authors, Timothy Higham at the University of California, Riverside, and Anthony Russell at the University of Calgary, Canada, both zoologists, brought Niklas into the project for his expertise on plant biomechanics.

"The paper shows that [adhesive capability] might be exaggerated, because geckos experience falls and a necessity to grip a surface like a leaf that requires a much more tenacious adhesion force; the paper shows that in some cases the adhesive ability can be exceeded," Niklas said.

In the theoretical study, the researchers developed computer models to understand if there are common-place instances when the geckos' ability to hold on to surfaces might be challenged, such as when canopy-dwelling geckos are being chased by a predator and are forced to leap from a tree, hoping to land on a leaf below. The researchers incorporated

ecological observations, adhesive force measurements, and body size and shape measurements of museum specimens to conduct simulations. They also considered the biomechanics of the leaves, the size of the leaves and the angles on the surface that geckos might land on to determine impact forces. Calculations were also based on worst-case scenarios, where a gecko reaches a maximum speed when it is no longer accelerating, called "terminal settling velocity."

"Leaves are cantilevered like diving boards and they go through harmonic motion [when struck], so you have to calculate the initial deflection and orientation, and then consider how does that leaf rebound and can the gecko still stay attached," Niklas said.

The final result showed that in some cases, geckos don't have enough adhesion to save themselves, he added.

Higham and Russell are planning to travel to French Guiana to do empirical adhesive force studies on living geckos in native forests.

The basic research helps people better understand how geckos stick to surfaces, and has the potential for future applications that mimic such biological mechanisms.

More information: Timothy E. Higham et al. Leaping lizards landing on leaves: escape-induced jumps in the rainforest canopy challenge the adhesive limits of geckos, *Journal of The Royal Society Interface* (2017). [DOI: 10.1098/rsif.2017.0156](https://doi.org/10.1098/rsif.2017.0156)

Provided by Cornell University

Citation: Gripping geckos' aerial escapes test their limits (2017, July 19) retrieved 30 September

2023 from <https://techxplore.com/news/2017-07-geckos-aerial-limits.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.