

Team improves optical efficiency in nanophotonic devices

January 4 2018, by Daniel Parry




Image shows directly measured polaritons propagating through a flake of Hexagonal boron nitride (hBN). This material has been identified as an ideal substrate for two-dimensional materials research while also recently being demonstrated as an exciting optical material for infrared nanophotonics. Credit: US Naval Research Laboratory

A team of physicists, headed by the U.S. Naval Research Laboratory (NRL), have demonstrated the means to improve the optical loss characteristics and transmission efficiency of hexagonal boron nitride devices, enabling very small lasers and nanoscale optics.

"The applications for this research are considerably broad," said Dr. Alexander J. Giles, research physicist, NRL Electronics Science and Technology Division. "By confining light to very small dimensions, nanophotonic devices have direct applications for use in ultra-high resolution microscopes, [solar energy harvesting](#), optical computing and targeted medical therapies."

Hexagonal boron nitride (hBN) forms an atomically thin lattice consisting of boron and nitrogen atoms. This material has recently been demonstrated as an exciting optical material for infrared nanophotonics and is considered an 'ideal substrate' for two-dimensional [materials](#).

While previous work demonstrated that natural hBN supports deeply sub-diffractive hyperbolic phonon polaritons desired for applications, such as, sub-diffractive optical imaging (so-called 'hyperlensing'), energy conversion, [chemical sensing](#), and quantum nanophotonics, limited transmission efficiencies continue to persist.

"We have demonstrated that the inherent efficiency limitations of nanophotonics can be overcome through the careful engineering of

isotopes in polar semiconductors and dielectric materials," Giles said.

Naturally occurring boron is comprised of two isotopes, boron-10 and boron-11, lending a 10 percent difference in atomic masses. This difference results in substantial losses due to phonon scattering, limiting the potential applications of this material. The research team at NRL has engineered greater than 99 percent isotopically pure samples of hBN, meaning they consist almost entirely of either boron-10 or boron-11 isotopes.

This approach results in a dramatic reduction in optical losses, resulting in optical modes that travel up to three times farther and persist for up to three times longer than natural hBN. These long-lived vibrational modes not only enable immediate advances specific to hBN - near field optics and chemical sensing - but also provide a strategic approach for other materials systems to exploit and build upon.

"Controlling and manipulating light at nanoscale, sub-diffractive dimensions is notoriously difficult and inefficient," said Giles. "Our work represents a new path forward for the next generation of materials and devices."

More information: Alexander J. Giles et al, Ultralow-loss polaritons in isotopically pure boron nitride, *Nature Materials* (2017). [DOI: 10.1038/nmat5047](https://doi.org/10.1038/nmat5047)

Provided by Naval Research Laboratory

Citation: Team improves optical efficiency in nanophotonic devices (2018, January 4) retrieved 13 July 2024 from <https://techxplore.com/news/2018-01-team-optical-efficiency-nanophotonic-devices.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.