

## Integrated modulator is now better than ever, made by the best optical materials

September 24 2018, by Leah Burrows



Conventional lithium niobate modulators [right] are the backbone of modern telecommunications, converting electronic data to optical information in fiber optic cables but are bulky, expensive and power hungry. This integrated, on-chip modulator [center] is 100 times smaller and 20 times more efficient. Credit: Harvard SEAS

Conventional lithium niobite modulators, the longtime workhorse of the



optoelectronic industry, may soon go the way of the vacuum tube and floppy disc. Researchers from the Harvard John A. Paulson School of Engineering and Applied Sciences have developed a new method to fabricate and design integrated, on-chip modulators 100 times smaller and 20 times more efficient than current lithium niobite (LN) modulators.

The research is described in *Nature*.

"This research demonstrates a fundamental technological breakthrough in integrated photonics," said Marko Loncar, the Tiantsai Lin Professor of Electrical Engineering at SEAS and senior author of the paper. "Our platform could lead to large-scale, very fast and ultra-low-loss photonic circuits, enabling a wide range of applications for future quantum and classical photonic communication and computation."

Harvard's Office of Technology Development (OTD) has worked closely with the Loncar Lab on the formation of a startup company, HyperLight, that intends to commercialize a portfolio of foundational intellectual property related to this research. Readying the technology toward the launch of HyperLight has been aided by funding from OTD's Physical Sciences & Engineering Accelerator, which provides translational funding for research projects that show potential for significant commercial impact.

Lithium niobate modulators are the backbone of modern telecommunications, converting electronic data to optical information in fiber optic cables. However, conventional LN modulators are bulky, expensive and power hungry. These modulators require a drive voltage of 3 to 5 volts, significantly higher than that provided by typical CMOS circuitry, which provides about 1 volt. As a result, separate, powerconsuming amplifiers are needed to drive the modulators, severely limiting chip-scale optoelectronic integration.







Fiber-optical networks, the backbone of the internet, rely on high-fidelity information conversion from electrical to the optical domain. The researchers combined the best optical material with innovative nanofabrication and design approaches, to realize, energy-efficient, high-speed, low-loss, electro-optic converters for quantum and classical communications. Credit: Second Bay Studios/Harvard SEAS

"We show that by integrating lithium niobate on a small chip, the drive voltage can be reduced to a CMOS-compatible level," said Cheng Wang, co-first author of the paper, former Ph.D. student and postdoctoral fellow at SEAS, and currently Assistant Professor at City University of Hong Kong. "Remarkably, these tiny modulators can also support data transmission rates up to 210 Gbit/s. It's like Antman—smaller, faster and better."

"Highly-integrated yet high-performance optical modulators are very important for the closer integration of optics and digital electronics, paving the way towards future fiber-in-fiber-out opto-electronic processing engines," said Peter Winzer, Director of Optical Transmission Research at Nokia Bell Labs, the industrial partner in this project, and coauthor of the paper. "We see this new <u>modulator</u> technology as a promising candidate for such solutions."

Lithium niobite is considered by many in the field to be difficult to work with on small scales, an obstacle that has so far ruled out practical integrated, on-chip applications. In previous research, Loncar and his team demonstrated a technique to fabricate high-performance lithium niobate microstructures using standard plasma etching to physically sculpt microresonators in thin lithium niobate films.



Combining that technique with specially designed electrical components, the researchers can now design and fabricate an integrated, highperformance on-chip modulator.

"Previously, if you wanted to make modulators smaller and more integrated, you had to compromise their performance," said Mian Zhang, a postdoctoral fellow at SEAS and co-first author of the research. "For example, existing integrated modulators can easily lose majority of the light as it propagates on the chip. In contrast, we have reduced losses by more than an order of magnitude. Essentially, we can control light without losing it."

"Because a modulator is such a fundamental component of communication technology—with a role equivalent to that of a transistor in computation technology—the applications are enormous," said Zhang. "The fact that these modulators can be integrated with other components on the same platform could provide practical solutions for nextgeneration long distance optical networks, data center optical interconnects, wireless communications, radar, sensing and so on."

**More information:** Cheng Wang et al, Integrated lithium niobate electro-optic modulators operating at CMOS-compatible voltages, *Nature* (2018). DOI: 10.1038/s41586-018-0551-y

Provided by Harvard John A. Paulson School of Engineering and Applied Sciences

Citation: Integrated modulator is now better than ever, made by the best optical materials (2018, September 24) retrieved 4 May 2024 from <u>https://techxplore.com/news/2018-09-modulator-optical-materials.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.