


Researchers advance organ-on-chip technology to advance drug development

August 23 2019



A 3D self-rolled biosensor array gripping a 3D cardiac spheroid. Credit: Carnegie Mellon University

Researchers from Carnegie Mellon University (CMU) and Nanyang Technological University, Singapore (NTU Singapore) have developed an organ-on-an-electronic-chip platform, which uses bioelectrical

sensors to measure the electrophysiology of the heart cells in three dimensions. These 3-D, self-rolling biosensor arrays coil up over heart cell spheroid tissues to form an "organ-on-e-chip," thus enabling the researchers to study how cells communicate with each other in multicellular systems such as the heart.

The organ-on-e-chip approach will help develop and assess the efficacy of drugs for disease treatment—perhaps even enabling researchers to screen for drugs and toxins directly on a human-like [tissue](#), rather than testing on animal tissue. The platform will also be used to shed light on the connection between the heart's [electrical signals](#) and disease, such as arrhythmias. The research, published in *Science Advances*, allows the researchers to investigate processes in cultured cells that currently are not accessible, such as tissue development and cell maturation.


"For decades, electrophysiology was done using cells and cultures on two-dimensional surfaces, such as culture dishes," says Associate Professor of Biomedical Engineering (BME) and Materials Science & Engineering (MSE) Tzahi Cohen-Karni. "We are trying to circumvent the challenge of reading the heart's electrical patterns in 3-D by developing a way to shrink-wrap sensors around heart cells and extracting electrophysiological information from this tissue."

The "organ-on-e-chip" platform starts out as a small, flat rectangle, not unlike a microscale slap bracelet. A slap bracelet starts out as a rigid, ruler-like structure, but when you release the tension it quickly coils up to band around the wrist.

The organ-on-e-chip starts out similarly. The researchers pin an array of sensors made of either metallic electrodes or graphene sensors to the chip's surface, then etch off a bottom layer of germanium, which is known as the "sacrificial layer." Once this sacrificial layer is removed, the biosensor array is released from its hold and rolls up from the

surface in a barrel shaped structure.

The researchers tested the platform on cardiac spheroids, or elongated organoids made of heart cells. These 3-D heart spheroids are about the width of 2-3 human hairs. Coiling the platform over the spheroid allows the researchers to collect electrical signal readings with high precision.




Calcium ion activity imaging of an encapsulated cardiac spheroid in the biosensor array. Credit: Carnegie Mellon University College of Engineering

"Essentially, we have created 3-D self-rolling biosensor arrays for exploring the electrophysiology of induced pluripotent stem cell derived cardiomyocytes," says lead author of the study and BME Ph.D. student Anna Kalmykov. "This platform could be used to do research into cardiac tissue regeneration and maturation that potentially can be used to

treat damaged tissue after a heart attack, for example, or developing new drugs to treat disease."

Through collaboration with the labs of BME/MSE Professor Adam Feinberg and former CMU faculty Jimmy Hsia, now Dean of the Graduate College of NTU Singapore, the researchers were able to design a proof of concept and test them on 3-D micro-mold formed cardiomyocyte spheroids.



Highly flexible 3D biosensor array unrolling with a glass capillary. Credit: Carnegie Mellon University College of Engineering

"Mechanics analysis of the roll-up process enables us to precisely control the shape of the sensors to ensure conforming contact between the sensors and the cardiac tissue," says NTU Professor Jimmy Hsia. "The technique also automatically adjusts the level of the delicate 'touch' between the sensors and the tissue such that high quality electric signals are measured without changing in the physiological conditions of the tissue due to external pressure."

"The whole idea is to take methods that are traditionally done in planar geometry and do them in three dimensions," says Cohen-Karni. "Our organs are 3-D in nature. For many years, electrophysiology was done using just [cells](#) cultured on a 2-D tissue culture dish. But now, these amazing electrophysiology techniques can be applied to 3-D structures."

More information: A. Kalmykov et al., "Organ-on-a-chip: Three-dimensional self-rolled biosensor array for electrical interrogations of human electrogenic spheroids," *Science Advances* (2019).

advances.sciencemag.org/content/5/8/eaax0729

Provided by Carnegie Mellon University

Citation: Researchers advance organ-on-chip technology to advance drug development (2019, August 23) retrieved 30 April 2024 from <https://techxplore.com/news/2019-08-advance-organ-on-chip-technology-drug.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
