


A new facial analysis method detects genetic syndromes with high precision and specificity

November 13 2019



Architecture of the proposed method for 3-D facial reconstruction and identification of facial dysmorphology associated with genetic syndromes. The green and red dots denote the correspondence of anatomical features between 2-D photographs and the statistical shape model (SSM) of the face. The classification uses both the geometry of the form (angles and distances) and the characteristics of texture (calculated around the locations of the yellow dots). Credit: Springer Nature Switzerland AG 2019.

Each year, over a million children are born with a genetic disease. Although about half of genetic syndromes present facial dysmorphology,

abnormal facial features are often subtle at birth and their identification by paediatricians can prove challenging. Delays and errors in diagnosis have a significant impact on mortality and morbidity associated with genetic syndromes. By way of example, the average accuracy in the detection of one of the most studied genetic syndromes, Down syndrome, by a trained paediatrician is as low as 64% in the US, and so methods for the early detection of genetic syndromes become very important.

Today, facial analysis of children from photographs is a technique that allows early identification of genetic syndromes. However, images may suffer problems of calibration and illumination. Although 3-D photography overcomes some of these problems, 3-D scanners to quantify craniofacial dysmorphology in children are expensive and often not available at all health centres. A recent study presents a new method to optimize facial analysis that enables reconstructing the face in 3-D from 2-D photographs.

Araceli Morales, Gemma Piella and Federico Sukno, members of the SIMBIOsys research group and of the Cognitive Media Technologies of the Department of Information and Communication Technologies (DTIC) at UPF, together with researchers from the University of Washington (USA) are the authors of this work published on 7 October in the online edition of *Lecture Notes in Computer Science*. The article describes the new optimization method to perform 3-D facial reconstructions of the shape of children's faces from uncalibrated 2-D photographs using a new statistical model.

First, for each 2-D photo, the new method estimates the camera pose using a statistical model and a set of 2-D facial landmarks. Secondly, the method calculates the camera pose and the parameters of the [statistical model](#) by minimizing the distance between the projection of the estimated 3-D face in the image plane of each camera and the observed

2-D face geometry.

"Using reconstructed 3-D faces, we automatically extract a set of 3-D geometric and appearance descriptors and we use them to train a classifier to identify facial dysmorphology associated with genetic syndromes," explains Araceli Morales, first author of the article who is working on this research for her doctoral thesis which is being supervised by Federico Sukno.

The face reconstruction method on 3-D photographs was evaluated in 54 subjects (age range 0-3 years), and "our classifier identified genetic syndromes in reconstructed 3-D faces from 2-D photographs with 100% sensitivity and a specificity of 92.11%," the authors explain in their article.

More information: Liyun Tu et al, Three-Dimensional Face Reconstruction from Uncalibrated Photographs: Application to Early Detection of Genetic Syndromes, *Uncertainty for Safe Utilization of Machine Learning in Medical Imaging and Clinical Image-Based Procedures* (2019). [DOI: 10.1007/978-3-030-32689-0_19](https://doi.org/10.1007/978-3-030-32689-0_19)

Provided by Universitat Pompeu Fabra - Barcelona

Citation: A new facial analysis method detects genetic syndromes with high precision and specificity (2019, November 13) retrieved 8 April 2024 from <https://techxplore.com/news/2019-11-facial-analysis-method-genetic-syndromes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.