

# Drones and AI detect soybean maturity with high accuracy

December 7 2020, by Lauren Quinn

---



Credit: Julio César García from Pixabay

Walking rows of soybeans in the mid-summer heat is an exhausting but essential chore in breeding new cultivars. Researchers brave the heat daily during crucial parts of the growing season to look for plants showing desirable traits, such as early pod maturity. But without a way to automate detection of these traits, breeders can't test as many plots as they'd like in a given year, elongating the time it takes to bring new cultivars to market.

In a new study from the University of Illinois, researchers predict soybean maturity date within two days using drone images and artificial intelligence, greatly reducing the need for boots on the ground.

"Assessing pod maturity is very time consuming and prone to errors. It's a [scoring system](#) based on the color of the pod, so it is also subject to human bias," says Nicolas Martin, assistant professor in the Department of Crop Sciences at Illinois and co-author on the study. "Many research groups are trying to use drone pictures to assess maturity, but can't do it at scale. So we came up with a more precise way to do that. It was really cool, actually."

Rodrigo Trevisan, a doctoral student working with Martin, trained computers to detect changes in canopy color from drone images collected across five trials, three growing seasons, and two countries. Importantly, he was able to account for "bad" images to maintain accuracy.

"Let's say we want to collect images every three days, but one day, there are clouds or it's raining, so we cannot. In the end, when you get the data from different years or different locations, they will all look different in terms of the number of images and the intervals and so on," Trevisan says. "The main innovation we developed is how we can account for whatever we are able to collect. Our model performs well independent of how often the data was collected."

Trevisan used a type of artificial intelligence called deep convolutional neural networks. He says CNNs are similar to the way human brains learn to interpret components of images—color, shape, texture—from our eyes.

"CNNs detect slight variations in color in addition to shapes, borders, and texture. For what we were trying to do, color was the most important

thing," Trevisan says. "But the advantage of the [artificial intelligence](#) models we used is that it would be quite straightforward to use the same model to predict another trait, such as yield or lodging. So now that we have these models set up, it should be much easier for people to use the same architecture and the same strategy to do many more things."

Martin says commercial breeding companies are clamoring for these capabilities.

"We had industry partners on the study who definitely want to use this in the years to come. And they made very good, important contributions. They wanted to make sure the answers were relevant for breeders in the field making decisions, selecting plants, and for farmers," Martin says. "Finding a good method to help breeders actually make decisions on large scales is quite exciting."

**More information:** Rodrigo Trevisan et al, High-Throughput Phenotyping of Soybean Maturity Using Time Series UAV Imagery and Convolutional Neural Networks, *Remote Sensing* (2020). [DOI: 10.3390/rs12213617](#)

Provided by University of Illinois at Urbana-Champaign

Citation: Drones and AI detect soybean maturity with high accuracy (2020, December 7) retrieved 9 April 2024 from <https://techxplore.com/news/2020-12-drones-ai-soybean-maturity-high.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---