

Engineers introduce the Oreometer

April 19 2022, by Jennifer Chu



Credit: Pixabay/CC0 Public Domain

When you twist open an Oreo cookie to get to the creamy center, you're mimicking a standard test in rheology—the study of how a non-Newtonian material flows when twisted, pressed, or otherwise stressed. MIT engineers have now subjected the sandwich cookie to rigorous materials tests to get to the center of a tantalizing question: Why does the cookie's cream stick to just one wafer when twisted apart?

"There's the fascinating problem of trying to get the cream to distribute evenly between the two wafers, which turns out to be really hard," says Max Fan, an undergraduate in MIT's Department of Mechanical Engineering.

In pursuit of an answer, the team subjected cookies to standard rheology tests in the lab and found that no matter the flavor or amount of stuffing, the cream at the center of an Oreo almost always sticks to one [wafer](#) when twisted open. Only for older boxes of cookies does the cream sometimes separate more evenly between both wafers.

The researchers also measured the torque required to twist open an Oreo, and found it to be similar to the torque required to turn a doorknob and about 1/10th what's needed to twist open a bottlecap. The cream's failure stress—i.e. the force per area required to get the cream to flow, or deform—is twice that of cream cheese and peanut butter, and about the same magnitude as mozzarella cheese. Judging from the cream's response to stress, the team classifies its texture as "mushy," rather than brittle, tough, or rubbery.

So, why does the cookie's cream glom to one side rather than splitting evenly between both? The manufacturing process may be to blame.

"Videos of the [manufacturing process](#) show that they put the first wafer down, then dispense a ball of cream onto that wafer before putting the second wafer on top," says Crystal Owens, an MIT mechanical engineering Ph.D. candidate who studies the properties of complex fluids. "Apparently that little time delay may make the cream stick better to the first wafer."

The team's study isn't simply a sweet diversion from bread-and-butter research; it's also an opportunity to make the science of rheology accessible to others. To that end, the researchers have designed a 3D-

printable "Oreometer"—a simple device that firmly grasps an Oreo cookie and uses pennies and rubber bands to control the twisting force that progressively twists the cookie open. Instructions for the tabletop device can be found [here](#).

The new study, "On Oreology, the fracture and flow of 'milk's favorite cookie," appears today in Kitchen Flows, a special issue of the journal *Physics of Fluids*. It was conceived of early in the COVID-19 pandemic, when many scientists' labs were closed or difficult to access. In addition to Owens and Fan, co-authors are mechanical engineering professors Gareth McKinley and A. John Hart.

Confection connection

A standard test in rheology places a fluid, slurry, or other flowable material onto the base of an instrument known as a rheometer. A parallel plate above the base can be lowered onto the test material. The plate is then twisted as sensors track the applied rotation and torque.

Owens, who regularly uses a laboratory rheometer to test fluid materials such as 3D-printable inks, couldn't help noting a similarity with sandwich cookies. As she writes in the new study:

"Scientifically, sandwich cookies present a paradigmatic model of parallel plate rheometry in which a fluid sample, the cream, is held between two parallel plates, the wafers. When the wafers are counter-rotated, the cream deforms, flows, and ultimately fractures, leading to separation of the cookie into two pieces."

While Oreo cream may not appear to possess fluid-like properties, it is considered a "yield stress fluid"—a soft solid when unperturbed that can start to flow under enough stress, the way toothpaste, frosting, certain cosmetics, and concrete do.

Curious as to whether others had explored the connection between Oreos and rheology, Owens found mention of a 2016 Princeton University study in which physicists first reported that indeed, when twisting Oreos by hand, the cream almost always came off on one wafer.

"We wanted to build on this to see what actually causes this effect and if we could control it if we mounted the Oreos carefully onto our rheometer," she says.

Cookie twist

In an experiment that they would repeat for multiple cookies of various fillings and flavors, the researchers glued an Oreo to both the top and bottom plates of a rheometer and applied varying degrees of torque and angular rotation, noting the values that successfully twisted each cookie apart. They plugged the measurements into equations to calculate the cream's viscoelasticity, or flowability. For each experiment, they also noted the cream's "post-mortem distribution," or where the cream ended up after twisting open.

In all, the team went through about 20 boxes of Oreos, including regular, Double Stuf, and Mega Stuf levels of filling, and regular, dark chocolate, and "golden" wafer flavors. Surprisingly, they found that no matter the amount of cream filling or flavor, the cream almost always separated onto one wafer.

"We had expected an effect based on size," Owens says. "If there was more cream between layers, it should be easier to deform. But that's not actually the case."

Curiously, when they mapped each cookie's result to its original position in the box, they noticed the cream tended to stick to the inward-facing wafer: Cookies on the left side of the box twisted such that the cream

ended up on the right wafer, whereas cookies on the right side separated with cream mostly on the left wafer. They suspect this box distribution may be a result of post-manufacturing environmental effects, such as heating or jostling that may cause cream to peel slightly away from the outer wafers, even before twisting.

The understanding gained from the properties of Oreo cream could potentially be applied to the design of other complex fluid materials.

"My 3D printing fluids are in the same class of materials as Oreo cream," she says. "So, this new understanding can help me better design ink when I'm trying to print flexible electronics from a slurry of carbon nanotubes, because they deform in almost exactly the same way."

As for the [cookie](#) itself, she suggests that if the inside of Oreo wafers were more textured, the cream might grip better onto both sides and split more evenly when twisted.

"As they are now, we found there's no trick to twisting that would split the [cream](#) evenly," Owens concludes.

More information: On Oreology, the fracture and flow of 'milk's favorite cookie', *Physics of Fluids* (2022). [DOI: 10.1063/5.0085362](https://doi.org/10.1063/5.0085362)

Provided by Massachusetts Institute of Technology

Citation: Engineers introduce the Oreometer (2022, April 19) retrieved 10 December 2023 from <https://techxplore.com/news/2022-04-oreometer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.