

Odorant analysis 2.0: Technique for the isolation of volatile food compounds optimized

July 29 2022, by Gisela Olias



Laboratory of the Research Group Food Metabolome Chemistry. Credit: J. Krpelan

A research team from the Leibniz Institute for Food Systems Biology at

the Technical University of Munich (LSB) has succeeded in automating an established method for the gentle, artifact-avoiding isolation of volatile food ingredients. As the team's current comparative study now shows, automated solvent-assisted flavor evaporation (aSAFE) offers significant advantages over the manual process. It achieves higher yields on average and reduces the risk of contamination by nonvolatile substances.

The optimized method is particularly important for odorant analysis. Odorants contribute significantly to the sensory profile of food and have a major influence on eating pleasure. Knowing the key odorants that shape the aroma of a food is therefore of interest both for analytical quality control and for targeted product development in the food industry.

Isolating volatile compounds from food is anything but trivial

However, isolating volatile compounds from food is not trivial. Many established methods lead to losses of labile odorants as well as to odor-active artifacts and are therefore unsuitable for [odorant](#) research. The manual SAFE technique developed in 1999 made it possible for the first time to easily isolate even thermally labile odorants from food without artifact formation. "This is an important prerequisite for using further [analytical methods](#) to identify the key odorants," says Philipp Schlumpberger, who contributed equally to the study with Christine Stübner. Both are currently working on their doctorates at LSB.

Today, manual SAFE is established worldwide as a standard procedure in aroma research. Nevertheless, the research team saw a need for optimization in ease of use, in the yields achieved, and in reducing the risk of transferring nonvolatile material, which can significantly

interfere with subsequent analytical steps.

The valve is critical

"As we discovered, the problems are mainly associated with the manual operation of the valve on the dropping funnel. Therefore, we replaced it with an electronically controlled pneumatic valve. To fully automate the SAFE apparatus, we optionally extended it with an automatic liquid nitrogen refill system as well as an endpoint detection and shutdown system," explains Martin Steinhaus, section and working group leader at LSB.

As the team's study now shows, the installation of the automatic valve increased yields, particularly for lipid-rich [food](#) extracts and for odorants with comparatively high boiling points. In addition, operator errors, which can lead to contamination of isolates with nonvolatile substances in the manual version, are eliminated with the automated SAFE.

"In the meantime, automated SAFE has replaced the manual variant in our laboratories. Other academic and industrial research groups are already following our example," says principal investigator Martin Steinhaus.

More information: Philipp Schlumpberger et al, Development and evaluation of an automated solvent-assisted flavour evaporation (aSAFE), *European Food Research and Technology* (2022). [DOI: 10.1007/s00217-022-04072-1](https://doi.org/10.1007/s00217-022-04072-1)

Provided by Leibniz-Institut für Lebensmittel-Systembiologie

Citation: Odorant analysis 2.0: Technique for the isolation of volatile food compounds optimized (2022, July 29) retrieved 16 April 2024 from <https://techxplore.com/news/2022-07-odorant-analysis-technique-isolation-volatile.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.