


# Preventing dye aggregation with molten salts to improve solar cell performance

August 29 2022



By modifying the surface of the oxide electrode with molten ionic salts, one can prevent the clumping of dye molecules without significantly impacting their coverage. This simple modification greatly improves the energy-conversion performance of dye-sensitized solar cells, find researchers from NITech, Japan in a new study. Credit: Tomohiko Inomata from NITech, Japan

Solar cells are quickly becoming one of the main ways to produce clean electricity in many countries in the world. Over the past few decades, a tremendous amount of effort has been dedicated to making solar power more prominent. However, the technology currently faces several challenges that limit widespread application.

In the case of dye-sensitized [solar cells](#) (DSSCs)—a highly promising photovoltaic technology—one of the main problems is dye aggregation. By design, DSSCs are electrochemical systems that mimic photosynthesis in plants; they rely on special photosensitive dyes to convert sunlight into electricity. Ideally, the dye should be applied evenly over the surface of an oxide electrode behind a transparent layer so that energy from absorbed sunlight can be transferred easily to the dye's electrons. This process generates free electrons that power an external circuit. However, most dyes tend to aggregate over the electrode surface in a way that hinders the desired flow of both light and electric charges. This takes a toll on the performance of DSSCs that has proven difficult to overcome.

Fortunately, a team of scientists led by Associate Professor Tomohiko Inomata of Nagoya Institute of Technology, Japan, may have just found a solution to this problem. In their recent study published in *RSC Advances*, they showed that certain ionic liquids ([molten salts](#) that are in [liquid state](#) at relatively low temperatures) can suppress dye aggregation to an impressive degree. Other members of this research team included Ms. Ayaka Matsunaga and Prof. Tomohiro Ozawa from Nagoya Institute of Technology, and Prof. Hideki Masuda from Aichi Institute of Technology, Japan.

But, how do ionic liquids achieve this feat? To shed light on the exact mechanism at play, the researchers focused on two ionic liquids with markedly different molecular sizes and two types of dyes. Both the ionic liquids had a similar molecular structure comprising an anchor that binds well to the electrode ([titanium dioxide](#),  $\text{TiO}_2$ ), a main polymer chain linking this anchor to a phosphor atom, and three additional short polymer chains protruding from the phosphor atom and away from the main "vertical" chain.

The researchers submerged the  $\text{TiO}_2$  electrodes in solutions with

different dye-to-ionic-liquid proportions and carefully analyzed how the different molecules adhered to them. After optimizing the synthesis procedure, they found that DSSCs made using the ionic liquid with a longer molecular structure had a remarkably better performance than their counterparts with non-modified oxide electrodes. "The spatially bulky molecular structure of ionic liquids acts as an effective anti-aggregation agent without significantly impacting the amount of dye adsorbed into the electrode," explains Dr. Inomata. "Most importantly, the introduction of the larger ionic liquid improves all the photovoltaic parameters of the DSSCs."

Needless to say, improving solar cell technology could give us an edge in the fight against the ongoing energy and climate crisis. Although ionic liquids are typically expensive, the way it is used by the team is, in fact, cost-effective. "Put simply, the idea is to apply ionic liquids only at the required part of the device—in this case, the electrode's surface," states Dr. Inomata.

The team believes that the widespread use of electrodes modified with ionic liquids could pave the way for highly functional yet affordable materials for solar cells and catalytic systems. Since the structure of [ionic liquids](#) can be tuned during their synthesis, they offer a much-needed versatility as anti-aggregation agents.

**More information:** Tomohiko Inomata et al, Improvements in photoelectric performance of dye-sensitised solar cells using ionic liquid-modified TiO<sub>2</sub> electrodes, *RSC Advances* (2022). [DOI: 10.1039/d2ra03230a](https://doi.org/10.1039/d2ra03230a)

Provided by Nagoya Institute of Technology

Citation: Preventing dye aggregation with molten salts to improve solar cell performance (2022, August 29) retrieved 16 April 2024 from <https://techxplore.com/news/2022-08-dye-aggregation-molten-salts-solar.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.