


Engineers improve electrochemical sensing by incorporating machine learning

November 25 2022, by Mary Fetzter



Graphical abstract. Credit: *Analytica Chimica Acta* (2022). DOI: 10.1016/j.aca.2022.340447

Combining machine learning with multimodal electrochemical sensing

can significantly improve the analytical performance of biosensors, according to new findings from a Penn State research team. These improvements may benefit noninvasive health monitoring, such as testing that involves saliva or sweat. The findings were published this month in *Analytica Chimica Acta*.

The researchers developed a novel analytical platform that enabled them to selectively measure multiple biomolecules using a single sensor, saving space and reducing complexity as compared to the usual route of using multi-sensor systems.

In particular, they showed that their sensor can simultaneously detect small quantities of [uric acid](#) and tyrosine—two important biomarkers associated with kidney and cardiovascular diseases, diabetes, [metabolic disorders](#), and neuropsychiatric and eating disorders—in sweat and saliva, making the developed method suitable for personalized health monitoring and intervention.

Many biomarkers have similar molecular structures or overlapping electrochemical signatures, making it difficult to detect them simultaneously. Leveraging machine learning for measuring multiple biomarkers can improve the accuracy and reliability of diagnostics and as a result improve patient outcomes, according to the researchers. Further, sensing using the same device saves resources and biological sample volumes needed for tests, which is critical with clinical samples with scarce amounts.


"We developed a new approach to improve the performance of electrochemical biosensors by combining machine learning with multimodal measurement," said Aida Ebrahimi, Thomas and Sheila Roell Early Career Assistant Professor of Electrical Engineering and assistant professor of biomedical engineering.

"Using our optimized machine learning architecture, we could detect biomolecules in amounts 100 times lower than what conventional sensing methods can do."

The researchers' methodology features a hardware/software system that enables them to automatically gather and process information based on a [machine learning model](#) that is trained to identify biomolecules in biological fluids such as saliva and sweat, which are common choices for noninvasive health monitoring.

"The machine learning-powered electrochemical diagnostic approach presented in this paper may find broader application in multiplexed biochemical sensing," said Vinay Kammarchedu, 2022-23 Milton and Albertha Langdon Memorial Graduate Fellow in Electrical Engineering at Penn State and first author on the paper.

"For example, this method can be extended to a variety of other molecules, including food and water toxins, drugs and neurochemicals that are challenging to detect simultaneously using conventional electrochemical methods."



Combining machine learning with multimodal electrochemical sensing can improve the analytical performance of biosensors and benefit non-invasive health monitoring, such as testing that involves sweat or saliva. Credit: Provided by Vinay Kammarchedu

In their ongoing work, the researchers are applying this approach on such neurochemicals, which are difficult to detect due to similarities in their molecular structure and overlapping electrochemical signatures.

"Our methodology successfully used one material to differentiate and distinguish four neurochemicals that are important in diseases like Parkinson's and Alzheimer's," Ebrahimi said.

"While this preliminary data is promising, we must work further to be able to detect the lower levels of these neurochemicals in biological samples such as saliva."

Beyond the specific results with the uric acid and tyrosine, the researchers are excited about the potential and versatility of the methodology.

"It is a new way of designing electrochemical diagnostic methods that may be applied to a variety of applications beyond biomedical systems," Ebrahimi said.

Combined with innovations in material and device engineering for sensor development, the researchers' analytical method may provide opportunities in pharmaceuticals, life science research, food screening, detection of environmental toxins and biodefense, where accurate and multiplexed testing or in-line monitoring is needed.

Conventionally, multiplexing is achieved by spectroscopic methods that rely on bulky and expensive equipment that is more suited for lab-based analysis. In the researchers' current prototype stage, the hardware is benchtop sized. They are working to make a smaller system that can be implemented for more than just health monitoring.

"Ultimately, we envision a handheld and field-deployable device that will be easier to use and more readily available than the current practices used in laboratory or clinical settings," Kammarchedu said.

More information: Vinay Kammarchedu et al, A machine learning-based multimodal electrochemical analytical device based on eMoSx-LIG for multiplexed detection of tyrosine and uric acid in sweat and saliva, *Analytica Chimica Acta* (2022). [DOI: 10.1016/j.aca.2022.340447](https://doi.org/10.1016/j.aca.2022.340447)

Provided by Pennsylvania State University

Citation: Engineers improve electrochemical sensing by incorporating machine learning (2022, November 25) retrieved 13 March 2024 from
<https://techxplore.com/news/2022-11-electrochemical-incorporating-machine.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.