


Speaking the same language: How artificial neurons mimic biological neurons

November 7 2022, by Christian Schneider



A biological neuron and the OAN. **a**, Simplified schematic of a biological neuron. Action potentials, the basic cell-to-cell communication events, are generated by rapid transmembrane ion exchanges through ion channels, and they propagate across the axon. In myelinated cells, alternate myelin/non-myelin domains (nodes of Ranvier) contribute to the fast and long-range action potential propagation. Biological neurons are immersed in an electrochemical environment, such as an aqueous electrolyte. This extracellular space is a

common reservoir containing various biological carriers for signaling and processing (ions, biomolecules and so on). Noise is also present in this environment. Ionic channels on the membrane endow neurons with ionic/molecular specificity and recognition. **b**, Circuit diagram of the OAN. The main part is an OEND that displays S-shaped negative differential resistance (S-NDR) phenomena and is responsive to ionic and biomolecular species common to biological environments. The OEND consists of two OECTs, namely, T_1 and T_2 , that are connected via the $R_1 = 5 \text{ k}\Omega$ and $R_1 = 10 \text{ k}\Omega$ resistors in a cascade-like configuration with feedback. The OAN is formed when the OEND is connected to an RC element ($R = 10 \text{ k}\Omega$; $C = 6 \text{ nF}$ to $10 \text{ }\mu\text{F}$) and voltage source V_{in} . Here V_{out} and I_{out} are the resulting output voltage and current, respectively, of the OEND under the influence of V_{in} . G, gate; S, source; D, drain. **c**, Schematic of the OECT that forms the OEND. The channel of the OECT consists of an organic mixed ionic–electronic conductor (OMIEC), such as PEDOT:PSS. **d**, Sensing mechanism in PEDOT:PSS. Ionic or polyatomic ions interact with PEDOT:PSS and modulate the doping level and hole conductivity of PEDOT:PSS, which results in a change in the OECT drain current and threshold voltage. Credit: *Nature Electronics* (2022). DOI: 10.1038/s41928-022-00859-y

Artificial intelligence has long been a hot topic: a computer algorithm "learns" by being taught by examples: What is "right" and what is "wrong." Unlike a computer algorithm, the human brain works with neurons—cells of the brain. These are trained and pass on signals to other neurons. This complex network of neurons and the connecting pathways, the synapses, controls our thoughts and actions.

Biological signals are much more diverse when compared with those in conventional computers. For instance, neurons in a biological neural network communicate with ions, biomolecules and neurotransmitters. More specifically, neurons communicate either chemically—by emitting the messenger substances such as neurotransmitters—or via [electrical impulses](#), so-called "action potentials" or "spikes".

Artificial neurons are a current area of research. Here, the efficient communication between the biology and electronics requires the realization of [artificial neurons](#) that emulate realistically the function of their biological counterparts. This means artificial neurons capable of processing the diversity of signals that exist in biology. Until now, most artificial neurons only emulate their biological counterparts electrically, without taking into account the wet biological environment that consists of ions, biomolecules and neurotransmitters.

Scientists led by Paschalis Gkoupidenis, group leader in Paul Blom's department at the Max Planck Institute for Polymer Research, have now tackled this problem and developed the first bio-realistic artificial neuron. This neuron can work in a biological environment and is able to produce diverse spiking dynamics that can be found in biology and can therefore communicate with their "real" biological counterparts.

For this purpose, the group of Gkoupidenis realized a non-linear element made of organic soft matter, something that also exists in biological [neurons](#). "Such [an] artificial element could be the key for bio-realistic neuroprosthetics that will speak the same language as biology, enabling the efficient restoration, replacement or even augmentation of the nervous system functions," says Gkoupidenis.

This allowed the researchers to develop for the first time a realistic artificial neuron that can communicate in diverse ways—chemically or via ionic charge carriers—in biological environments.

Their results have now been published in the journal *Nature Electronics*.

More information: Fabrizio Torricelli, An organic artificial spiking neuron for in situ neuromorphic sensing and biointerfacing, *Nature Electronics* (2022). [DOI: 10.1038/s41928-022-00859-y](https://doi.org/10.1038/s41928-022-00859-y).
www.nature.com/articles/s41928-022-00859-y

Provided by Max Planck Society

Citation: Speaking the same language: How artificial neurons mimic biological neurons (2022, November 7) retrieved 27 April 2024 from <https://techxplore.com/news/2022-11-language-artificial-neurons-mimic-biological.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.