

Machine learning helps researchers separate compostable from conventional plastic waste with 'very high' accuracy

March 14 2023



Credit: Unsplash/CC0 Public Domain

Disposable plastics are everywhere: Food containers, coffee cups, plastic bags. Some of these plastics, called compostable plastics, can be

engineered to biodegrade under controlled conditions. However, they often look identical to conventional plastics, are recycled incorrectly, and as a result, contaminate plastic waste streams and reduce recycling efficiency. Similarly, recyclable plastics are often mistaken for compostable ones, resulting in polluted compost.

Researchers at University College London (UCL) have published a paper in *Frontiers in Sustainability* in which they used machine learning to automatically sort different types of [compostable](#) and [biodegradable plastics](#) and differentiate them from conventional plastics.

"The [accuracy](#) is very high and allows the technique to be feasibly used in industrial recycling and composting facilities in the future," said Prof Mark Miodownik, corresponding author of the study.

Up to perfect accuracy

The researchers worked with different types of plastics measuring between 50mm by 50mm and 5mm by 5mm. Conventional [plastic](#) samples included PP and PET, often used for [food containers](#) and drinking bottles, as well as LDPE, used—among other things—for [plastic bags](#) and packaging. Compostable plastic samples included PLA and PBAT, used for cup lids, tea bags, and magazine wraps; as well as palm leaf and sugarcane, both biomass-derived materials used to produce packaging. The samples were divided into a training set, used to build classification models; and a testing set, used to check accuracy.

Results showed high success rates: The model achieved perfect accuracy for all materials when the samples measured more than 10mm by 10mm. For sugarcane-derived or palm leaf-based materials measuring 10mm by 10mm or less, however, the misclassification rate was 20% and 40%, respectively.

Looking at pieces measuring 5mm by 5mm, some materials were identified more reliably than others: For LDPE and PBAT pieces the misclassification rate was 20%; and both biomass-derived materials were misidentified at rates of 60% (sugarcane) and 80% (palm leaf). The model was, however, able to identify PLA, PP and PET pieces without error, regardless of sample measurements.

Beyond the visible

"Currently, most compostable plastics are treated as a contaminant in the recycling of conventional plastics, reducing their value. Trommel and density sorting are applied to screen compost and reduce the presence of other materials. However, the level of contaminants from the current screening process is unacceptably high," explained Miodownik. "The advantages of compostable packaging are only realized when they are industrially composted and do not enter the environment or pollute other waste streams or the soil."

To improve accuracy, a team of scientists including Nutchaneepanichskul, Prof Helen Hailes, and Miodownik from UCL's Plastic Waste Innovation Hub tested different types of conventional, compostable, and biodegradable plastics, using [hyperspectral imaging](#) (HSI) for classification model development. HSI is an imaging technique that detects the invisible chemical signature of different materials while scanning them, producing a pixel-by-pixel chemical description of a sample. AI models were used to interpret these descriptions and make a material identification.

Plastic mismanagement in recycling and industrial composting processes is high, making reliable sorting mechanisms essential. "Currently, the speed of identification is too low for implementation at industrial scale," Miodownik pointed out. However, "we can and will improve it since automatic sorting is a key technology to make compostable [plastics](#) a

sustainable alternative to recycling."

More information: Machine learning helps researchers separate compostable from conventional plastic waste with 'very high' accuracy, *Frontiers in Sustainability* (2023). [DOI: 10.3389/frsus.2023.1125954](https://doi.org/10.3389/frsus.2023.1125954) , www.frontiersin.org/articles/10.3389/frsus.2023.1125954/full

Provided by Frontiers

Citation: Machine learning helps researchers separate compostable from conventional plastic waste with 'very high' accuracy (2023, March 14) retrieved 23 April 2024 from <https://techxplore.com/news/2023-03-machine-compostable-conventional-plastic-high.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.