

Researchers successfully train a machine learning model in outer space for the first time

July 28 2023



The ION SCV004 satellite, on which the machine learning models were run. The image shows the satellite releasing CubeSats (small satellites) into space. Credit: D-Orbit

For the first time, researchers have trained a machine learning model in outer space, on board a satellite. This achievement could enable real-time monitoring and decision making for a range of applications, from disaster management to deforestation.

This project has been summarized in a publication, "Fast model inference and training on-board of Satellites," available on the pre-print server *arXiv*. The work was also presented at the International Geoscience and Remote Sensing Symposium (IGARSS) conference on 21 July 2023.

Data collected by remote-sensing satellites is fundamental for many key activities, including aerial mapping, weather prediction, and monitoring deforestation. Currently, most satellites can only passively collect data, since they are not equipped to make decisions or detect changes. Instead, data has to be relayed to Earth to be processed, which typically takes several hours or even days. This limits the ability to identify and respond to rapidly emerging events, such as a natural disaster.

To overcome these restrictions, a group of researchers led by DPhil student Vít Růžička (Department of Computer Science, University of Oxford), took on the challenge of training the first machine learning program in [outer space](#).

During 2022, the team successfully pitched their idea to the Dashing through the Stars mission, which had issued an open call for project proposals to be carried out on board the ION SCV004 satellite, launched in January 2022. During the autumn of 2022, the team uplinked the code for the program to the satellite already in orbit.

The researchers trained a simple model to detect changes in [cloud cover](#) from aerial images directly onboard the satellite, in contrast to training on the ground. The model was based on an approach called few-shot

learning, which enables a model to learn the most important features to look for when it has only a few samples to train from. A key advantage is that the data can be compressed into smaller representations, making the model faster and more efficient.

Vít Ružička said, "The model we developed, called RaVAEn, first compresses the large image files into vectors of 128 numbers. During the training phase, the model learns to keep only the informative values in this vector; the ones that relate to the change it is trying to detect—in this case, whether there is a cloud present or not. This results in extremely fast training due to having only a very small classification model to train."

While the first part of the model, to compress the newly-seen images, was trained on the ground, the second part (which decided whether the image contained clouds or not) was trained directly on the satellite.

Normally, developing a [machine learning model](#) would require several rounds of training, using the power of a cluster of linked computers. In contrast, the team's tiny model completed the training phase (using over 1,300 images) in around one and a half seconds.

When the team tested the model's performance on novel data, it automatically detected whether a cloud was present or not in around a tenth of a second. This involved encoding and analyzing a scene equivalent to an area of about 4.8 x 4.8 km² area (equivalent to almost 450 football pitches).

According to the researchers, the model could easily be adapted to carry out different tasks, and to use other forms of data. Vít Ružička added, "Having achieved this demonstration, we now intend to develop more advanced models that can automatically differentiate between changes of interest (for instance flooding, fires, and deforestation) and natural

changes (such as natural changes in leaf color across the seasons)."

"Another aim is to develop models for more complex data, including images from hyperspectral satellites. This could allow, for instance, the detection of methane leaks, and would have key implications for combatting climate change."

Performing machine learning in outer space could also help overcome the problem of on-board satellite sensors being affected by the harsh environmental conditions, so that they require regular calibration.

Vít Růžička said, "Our proposed system could be used in constellations of non-homogeneous satellites, where reliable information from one [satellite](#) can be applied to train the rest of the constellation. This could be used, for instance, to recalibrate sensors that have degraded over time or experienced rapid changes in the environment."

More information: Vít Růžička et al, Fast model inference and training on-board of Satellites, *arXiv* (2023). [DOI: 10.48550/arxiv.2307.08700](#)

Provided by University of Oxford

Citation: Researchers successfully train a machine learning model in outer space for the first time (2023, July 28) retrieved 23 February 2024 from <https://techxplore.com/news/2023-07-successfully-machine-outer-space.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.