

## Inverted perovskite solar cell breaks 25% efficiency record

November 18 2023



Northwestern University researchers have raised the standards again for perovskite solar cells with a new development that helped the emerging technology hit new records for efficiency. Credit: Sargent Lab/Northwestern University

Northwestern University researchers have raised the standards again for



perovskite solar cells with a new development that helped the emerging technology hit new records for efficiency.

The findings, <u>published today (Nov. 17) in the journal *Science*, describe a dual-molecule solution to overcoming losses in efficiency as sunlight is converted to energy. By incorporating first, a molecule to address something called surface <u>recombination</u>, in which electrons are lost when they are trapped by defects—missing atoms on the surface, and a second molecule to disrupt recombination at the interface between layers, the team achieved a National Renewable Energy Lab (NREL) certified efficiency of 25.1% where earlier approaches reached efficiencies of just 24.09%.</u>

"Perovskite solar technology is moving fast, and the emphasis of research and development is shifting from the bulk absorber to the interfaces," said Northwestern professor Ted Sargent. "This is the <u>critical point</u> to further improve efficiency and stability and bring us closer to this promising route to ever-more-efficient solar harvesting."

Sargent is the co-executive director of the Paula M. Trienens Institute for Sustainability and Energy (formerly ISEN) and a multidisciplinary researcher in <u>materials chemistry</u> and <u>energy systems</u>, with appointments in the department of chemistry in the Weinberg College of Arts and Sciences and the department of electrical and computer engineering in the McCormick School of Engineering.

Conventional solar cells are made of high-purity silicon wafers that are energy-intensive to produce and can only absorb a fixed range of the solar spectrum.

Perovskite materials whose size and composition can be adjusted to "tune" the wavelengths of light they absorb, making them a favorable and potentially lower-cost, high-efficiency emerging tandem technology.



Historically <u>perovskite</u> solar cells have been plagued by challenges to improve efficiency because of their relative instability. Over the past few years, advances from Sargent's lab and others have brought the efficiency of <u>perovskite solar cells</u> to within the same range as what is achievable with silicon.

In the present research, rather than trying to help the cell absorb more sunlight, the team focused on the issue of maintaining and retaining generated electrons to increase efficiency. When the perovskite layer contacts the electron transport layer of the cell, electrons move from one to the other. But the electron can move back outward and fill, or "recombine" with holes that exist on the perovskite layer.

"Recombination at the interface is complex," said first author Cheng Liu, a postdoctoral student in the Sargent lab, which is co-supervised by the Charles E. and Emma H. Morrison Professor of Chemistry Mercouri Kanatzidis. "It's very difficult to use one type of molecule to address complex recombination and retain electrons, so we considered what combination of <u>molecules</u> we could use to more comprehensively solve the problem."

Past research from Sargent's team has found evidence that one molecule, PDAI<sub>2</sub>, does a good job at solving interface recombination. Next they needed to find a molecule that would work to repair surface defects and prevent electrons from recombining with them.

By finding the mechanism that would allow  $PDAI_2$  to work with a secondary molecule, the team narrowed in on sulfur, which could replace carbon groups—typically poor at preventing electrons from moving—to cover missing atoms and suppress recombination.

A recent paper by the same group published in *Nature* developed a coating for the substrate beneath the perovskite layer to help the cell



work at a higher temperature for a longer period. This solution, according to Liu, can work in tandem with the findings within the *Science* paper.

While the team hopes their findings will encourage the larger scientific community to continue moving the work forward, they too will be working on follow-ups.

"We have to use a more flexible strategy to solve the complex interface problem," Cheng said. "We can't only use one kind of molecule, as people previously did. We use two molecules to solve two kinds of recombination, but we are sure there's more kinds of defect-related recombination at the interface. We need to try to use more molecules to come together and make sure all molecules work together without destroying each other's functions."

**More information:** Cheng Liu et al, Bimolecularly passivated interface enables efficient and stable inverted perovskite solar cells, *Science* (2023). DOI: 10.1126/science.adk1633

Provided by Northwestern University

Citation: Inverted perovskite solar cell breaks 25% efficiency record (2023, November 18) retrieved 8 May 2024 from https://techxplore.com/news/2023-11-inverted-perovskite-solar-cell-efficiency.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.