


New design for a rechargeable hydrogen-chlorine battery in a wide temperature range

November 22 2023, by Liu Jia



Credit: *Journal of the American Chemical Society* (2023). DOI: 10.1021/jacs.3c09819

A research team led by Prof. Chen Wei from the University of Science and Technology of China (USTC) of the Chinese Academy of Sciences (CAS) designed a rechargeable hydrogen-chlorine ($\text{H}_2\text{-Cl}_2$) battery that can operate in temperatures ranging from -70°C to 40°C . The study was

[published](#) in *Journal of the American Chemical Society* as the cover article.

Hydrogen fuel cells are a promising technology valuable for their sustainability and the abundance of hydrogen, among which $\text{H}_2\text{-Cl}_2$ fuel cells stand out due to fast electrochemical kinetics, high redox potential and high specific capacity of Cl_2/Cl^- redox couple. However, the volatile chlorine gas cannot be retained during the charging process, resulting in poor Coulombic efficiency (CE) and reversibility. There is an urgent need to develop aqueous chlorine batteries with [high performance](#) and applicability at different temperatures.

The research team first discovered that due to the lack of binding sites with strong affinity to Cl_2 , traditional adsorptive cathodes have difficulty immobilizing Cl_2 , causing low reversibility. To tackle this problem, the team designed a hierarchically porous carbon [cathode](#) composed of highly micro-/mesoporous carbon (HPC) and macroporous carbon felt (CF), effectively confining the Cl_2 on the cathode and improving the reversibility.

The team showed that the $\text{H}_2\text{-Cl}_2$ cells maintained a high CE and stability, operating steadily for 500 cycles at a discharge capacity of 3 mAh cm^{-2} . In addition, the cells operate well at ultralow temperatures, maintaining a discharge plateau of 1.1 V and a high specific capacity of 282 mAh g^{-1} at -70°C .

To further understand the mechanism of reversibility improvement, the team combined X-ray photoelectron spectroscopy (XPS) with theoretical calculations and revealed that the Cl_2/Cl^- reaction occurs along with the reversible formation and breakage of C-Cl bonds, which enhances the [reversibility](#) of the Cl_2/Cl^- cathodes.

This work provides a new direction for the design of aqueous chlorine

batteries and high-energy-density hydrogen batteries in a wide temperature range.

More information: Zehui Xie et al, Rechargeable Hydrogen–Chlorine Battery Operates in a Wide Temperature Range, *Journal of the American Chemical Society* (2023). [DOI: 10.1021/jacs.3c09819](https://doi.org/10.1021/jacs.3c09819)

Provided by Chinese Academy of Sciences

Citation: New design for a rechargeable hydrogen-chlorine battery in a wide temperature range (2023, November 22) retrieved 12 May 2024 from <https://techxplore.com/news/2023-11-rechargeable-hydrogen-chlorine-battery-wide-temperature.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--