


# Electrolyte solution reduces Li-ion battery recharging time while capacity remains at low temperatures

March 1 2024, by Bob Yirka



Electrolyte design and solvent-screening strategies. **a–c**, Schematic illustrations of Li<sup>+</sup> transport behaviors with the vehicular mechanism (**a**), structural mechanism (**b**) and ligand-channel-facilitated mechanism (**c**). **d**, The solvation

energy and solvation-shell volume of solvents. e, The  $\text{Li}^+$  transport energy barrier and solvation-shell volume of solvents. The electrolyte using the solvents with small size, low solvation energy and  $\text{Li}^+$  transport energy barrier can conduct  $\text{Li}^+$  through the ligand-channel transport mechanism. Credit: *Nature* (2024). DOI: 10.1038/s41586-024-07045-4

A team of chemists and engineers affiliated with several institutions has found an electrolyte solution that can be used to reduce the recharging time of lithium-ion batteries while allowing battery capacity to remain comparatively high at low temperatures.

In their paper [published](#) in the journal *Nature*, the group describes the new electrolyte and how well it worked during testing. Chong Yan and Jia-Qi Huang with the Beijing Institute of Technology have published a [News and Views piece](#) in the same journal issue, outlining the work done by the team on this new effort.

As Yan and Huang note, [lithium-ion batteries](#) have proven their usefulness in a wide variety of applications, but that does not mean there is no room for improvement. One improvement that users of battery-powered devices would like to see is faster recharging. They would also like to see improvements in [battery capacity](#) as temperatures drop during the winter months. In this new study, the research team found an [electrolyte solution](#) that they claim reduces charging time while also preventing loss of capacity when exposed to [cold temperatures](#).

In their work, the team working in China found that the use of organic solvents could greatly improve the mobility of ions in a battery electrolyte, allowing for faster charging. They noted that such solvents could also be used to prevent loss of capacity in temperatures as low as  $-80^{\circ}\text{C}$ .

In their work, the team used a solvent called fluoroacetonitrile, which has molecules that are much smaller than those typically used to make the electrolyte. The molecules of the solvent tend to surround the lithium ions, forming a shell. As the shells touch, they form a sort of tunnel, allowing the ions to move more quickly through the electrolyte.

Testing showed ionic conductivity of 40.3 millisiemens per centimeter under normal ambient conditions, which, the team notes, is approximately four times that of standard batteries. They also found that putting their test batteries in a freezer did not reduce their capacity.

**More information:** Di Lu et al, Ligand-channel-enabled ultrafast Li-ion conduction, *Nature* (2024). [DOI: 10.1038/s41586-024-07045-4](https://doi.org/10.1038/s41586-024-07045-4)

Chong Yan et al, Tiny sheaths of solvent boost battery performance, *Nature* (2024). [DOI: 10.1038/d41586-024-00378-0](https://doi.org/10.1038/d41586-024-00378-0)

© 2024 Science X Network

Citation: Electrolyte solution reduces Li-ion battery recharging time while capacity remains at low temperatures (2024, March 1) retrieved 8 May 2024 from <https://techxplore.com/news/2024-03-electrolyte-solution-li-ion-battery.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--