

Trotting robots reveal emergence of animal gait transitions

April 30 2024, by Celia Luterbacher



Guillaume Bellegarda, Milad Shafiee and Auke Ijspeert. Credit: EPFL/Jamani Caillet—CC-BY-SA 4.0

A four-legged robot trained with machine learning by EPFL researchers has learned to avoid falls by spontaneously switching between walking, trotting, and pronking—a milestone for roboticists as well as biologists interested in animal locomotion.



With the help of a form of machine learning called <u>deep reinforcement</u> <u>learning</u> (DRL), the EPFL robot notably learned to transition from trotting to pronking—a leaping, arch-backed gait used by animals like springbok and gazelles—to navigate a challenging <u>terrain</u> with gaps ranging from 14 to 30 cm. The study, led by the BioRobotics Laboratory in EPFL's School of Engineering, offers new insights into why and how such gait transitions occur in animals.

"Previous research has introduced energy efficiency and musculoskeletal injury avoidance as the two main explanations for gait transitions. More recently, biologists have argued that stability on flat terrain could be more important.

"But animal and robotic experiments have shown that these hypotheses are not always valid, especially on uneven ground," says Ph.D. student Milad Shafiee, first author on <u>a paper</u> published in *Nature Communications*.

Shafiee and co-authors Guillaume Bellegarda and BioRobotics Lab head Auke Ijspeert were therefore interested in a new hypothesis for why gait transitions occur: viability, or fall avoidance. To test this hypothesis, they used DRL to train a quadruped robot to cross various terrains.

On flat terrain, they found that different gaits showed different levels of robustness against random pushes, and that the robot switched from a walk to a trot to maintain viability, just as quadruped animals do when they accelerate.

When confronted with successive gaps in the experimental surface, the robot spontaneously switched from trotting to pronking to avoid falls. Moreover, viability was the only factor that was improved by such gait transitions.



"We showed that on flat terrain and challenging discrete terrain, viability leads to the emergence of gait transitions, but that energy efficiency is not necessarily improved," Shafiee explains. "It seems that energy efficiency, which was previously thought to be a driver of such transitions, may be more of a consequence. When an animal is navigating challenging terrain, it's likely that its first priority is not falling, followed by energy efficiency."

A bio-inspired learning architecture

To model locomotion control in their robot, the researchers considered the three interacting elements that drive animal movement: the brain, the <u>spinal cord</u>, and sensory feedback from the body. They used DRL to train a <u>neural network</u> to imitate the spinal cord's transmission of brain signals to the body as the robot crossed an experimental terrain.

Then, the team assigned different weights to three possible learning goals: <u>energy efficiency</u>, force reduction, and viability. A series of computer simulations revealed that of these three goals, viability was the only one that prompted the robot to automatically—without instruction from the scientists—change its gait.

The team emphasizes that these observations represent the first learningbased locomotion framework in which <u>gait</u> transitions emerge spontaneously during the learning process, as well as the most dynamic crossing of such large consecutive gaps for a quadrupedal robot.

"Our bio-inspired learning architecture demonstrated state-of-the-art quadruped <u>robot</u> agility on the challenging terrain," Shafiee says.

The researchers aim to expand on their work with additional experiments that place different types of robots in a wider variety of challenging environments. In addition to further elucidating <u>animal</u>



<u>locomotion</u>, they hope that ultimately, their work will enable the more widespread use of robots for biological research, reducing reliance on animal models and the associated ethics concerns.

More information: Milad Shafiee et al, Viability leads to the emergence of gait transitions in learning agile quadrupedal locomotion on challenging terrains, *Nature Communications* (2024). DOI: 10.1038/s41467-024-47443-w

Provided by Ecole Polytechnique Federale de Lausanne

Citation: Trotting robots reveal emergence of animal gait transitions (2024, April 30) retrieved 17 July 2024 from <u>https://techxplore.com/news/2024-04-robots-reveal-emergence-animal-gait.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.