


# Scientists convert chicken fat into energy storage devices

May 10 2024



Graphical abstract. Credit: *ACS Applied Materials & Interfaces* (2024). DOI: 10.1021/acsami.4c02753

The global move toward more sustainable, green energy has increased power reserves and the demand for energy storage devices.

Unfortunately, some materials for these devices can be expensive and environmentally problematic. Producing alternative energy storage devices from things that are usually thrown away could help resolve these challenges.

Now, researchers in *ACS Applied Materials & Interfaces* [report](#) a method to transform [chicken](#) fat into carbon-based electrodes for supercapacitors that store energy and power LEDs.

In 2023, global renewable energy capacity experienced an unprecedented almost 50 percent increase versus the previous year, according to the International Energy Agency. But that excess energy must be stored somewhere for the world to benefit from its production later. For example, sunny days in California have recently triggered negative energy prices due to excess supply from rooftop solar panels.

Recent efforts to design high-performance storage devices have taken advantage of carbon materials, such as graphene, because of their efficient charge transportation and natural abundance, but their fabrication is expensive and generates pollution and [greenhouse gases](#).

Looking for an alternative carbon source material, Mohan Reddy Pallavolu, Jae Hak Jung, Sang Woo Joo, and colleagues wanted to develop a simple, cost-effective method for converting waste chicken fat into electrically conductive nanostructures for supercapacitor [energy storage devices](#).

The researchers first used a gas flame gun to render the fat from a chicken and burned the melted oil using a flame wick method, much as one would use an oil lamp. They then collected the soot on the bottom of a flask, which was suspended above the flame.

Electron microscopy showed that the soot contained carbon-based nanostructures that were uniform spherical lattices of concentric graphite rings, like the layers of onions. The researchers tested a way to enhance the electrical characteristics of the carbon nanoparticles by soaking them in a solution of thiourea.

Assembled into the negative electrode of an asymmetric supercapacitor, the chicken fat-sourced carbon nanoparticles demonstrated good capacitance and durability, as well as high energy and power density. As predicted, these properties were improved further when the electrodes were made of the thiourea-treated carbon nanoparticles.

The researchers then demonstrated that the new supercapacitor could perform real-time applications—charging and connecting two of them to light up red, green, and blue LEDs. The results highlight the potential advantages of using [food waste](#) like chicken fat as a [carbon source](#) in the search for even greener [green energy](#).

**More information:** Jyothi Nallapureddy et al, Strategic Way of Synthesizing Heteroatom-Doped Carbon Nano-onions Using Waste Chicken Fat Oil for Energy Storage Devices, *ACS Applied Materials & Interfaces* (2024). [DOI: 10.1021/acsami.4c02753](https://doi.org/10.1021/acsami.4c02753)

Provided by American Chemical Society

Citation: Scientists convert chicken fat into energy storage devices (2024, May 10) retrieved 13 June 2024 from <https://techxplore.com/news/2024-05-scientists-chicken-fat-energy-storage.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.