

Engineers study hovering bats and hummingbirds in Costa Rica

26 September 2018



A Purple-crowned fairy hovers at the feeder while instantaneous vertical forces are measured. Credit: Lentink Lab, Stanford University

Each sunrise in Las Cruces, Costa Rica, River Ingersoll's field team trekked into the jungle to put the finishing touches on nearly invisible nets. A graduate student in the lab of David Lentink, assistant professor of mechanical engineering at Stanford University, Ingersoll needed these delicate nets to catch, study and release the region's abundant hummingbirds and bats—the only two vertebrates with the ability to hover in place.

"We're really interested in how hovering flight evolved," said Ingersoll. "Nectar bats drink from flowers like hummingbirds do, so we want to see if there's any similarities or differences between these two different taxa."

Ingersoll's nets worked, and he ended up examining over 100 individual hummingbirds and bats, covering 17 hummingbird and three bat species, during his field study, the results of which the group published in *Science Advances*.

Through a combination of high-speed camera footage and aerodynamic force measurements, he and his fellow researchers found that hummingbirds and bats hover in very different

ways. Yet they also found that nectar bats' hovering shares some similarities with hummingbird hovering—which fruit bats do not share. This suggests that they evolved a different method to hover compared with other bats in order to drink nectar.


In addition to learning more about bats and hummingbirds, Lentink and others can apply what they learned to engineering problems, such as designing flying robots. Engineers have already created robots inspired by hummingbirds and bats but haven't known which of the natural counterparts of these robots hover most effectively.

Watching every feather

It is simple to imagine how a flying animal supports itself by flapping downward, but in order to avoid exaggerated bobbing up and down, hovering animals must maintain this support while flapping upward as well. Hummingbirds and bats accomplish this feat by twisting their wings backward on the upstroke, continuously pushing air downward to keep them steadily aloft.

"If you look amongst vertebrates, there are two that can hover in a sustained way," said Lentink. "Those are hummingbirds and nectar bats. And you'll find both in the neotropics, like Costa Rica."

To study these subjects, Ingersoll collaborated with a long-standing bird banding project run by Stanford ecologists in Las Cruces. Borrowing birds and bats from their project, he placed each animal in a flight chamber outfitted with aerodynamic force sensors at the top and bottom of the chamber—equipment developed by Lentink's lab to measure extremely small changes in vertical force at 10,000 times per second. These plates are so sensitive that they captured the vertical forces produced by every twist and flutter of hummingbirds that weighed as little as 2.4 grams.



A Garden Emerald hovers at the feeder while instantaneous vertical forces are measured. Credit: Lentink Lab, Stanford University

By synching those force measurements with multiple high-speed cameras recording at 2,000 frames per second, the researchers could isolate any moment of their subjects' flights to see how the lift they were generating related to the shape of their wings.

"I'd sit and wait for the hummingbird to feed at the flower. Once it was feeding, I would trigger the cameras and the force measurements and we'd get four seconds of footage of the [hummingbird](#) flapping at the flower," said Ingersoll.

After their short stint in the flight chamber, Ingersoll returned the birds and bats to where they were caught and released them. The whole process took between one and three hours.

Different ways to hover

The researchers found that the bats and hummingbirds all exerted a similar amount of energy relative to their weight during these flights but that the hummingbirds, fruit bats and nectar bats all hovered in very different ways. The hummingbirds hovered in a more aerodynamically efficient way than the bats—the hummingbirds generated more lift relative to drag. In comparing [wing](#) shapes, the researchers found this efficiency

is likely because the hummingbirds invert their wings more easily. Although the bats struggled with turning over their wings, they exerted a comparable amount of energy because they have bigger wings and larger strokes.

The researchers were surprised to find that nectar bats, which sidle up to flowers like hummingbirds, generated more upward force when the wings were lifting than fruit bats. Looking at their wing shape, the researchers found that nectar bats can twist their wings much more than fruit bats on the upstroke. So nectar bats' hovering form is like a blend of [fruit bats'](#) and hummingbirds' hovering.

The researchers plan to build on these findings as part of their work on flapping robots and drones but Lentink also sees potential for more work beyond the lab.

"When Rivers proposed to do this study in Costa Rica, a field study was something I'd never hoped for. Now, he really inspired me," said Lentink. "There are about 10,000 species of birds and most of them have never been studied. It sounds like too big a study to embark on but that's what I dream about."

More information: R. Ingersoll et al., "Biomechanics of hover performance in Neotropical hummingbirds versus bats," *Science Advances* (2018). advances.sciencemag.org/content/4/9/eaat2980

Provided by Stanford University

APA citation: Engineers study hovering bats and hummingbirds in Costa Rica (2018, September 26)
retrieved 17 January 2022 from <https://techxplore.com/news/2018-09-hummingbirds-costa-rica.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.